



## Contamination of lettuce with antibiotic resistant E. coli after slurry application

Jensen, Annette Nygaard; Storm, Christina; Baggesen, Dorte Lau; Forslund, A.; Dalsgaard, A.

*Published in:*

Book of Abstracts. First Int. Conference on Organic Food Quality and Health research

*Publication date:*

2011

*Document Version*

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Jensen, A. N., Storm, C., Baggesen, D. L., Forslund, A., & Dalsgaard, A. (2011). Contamination of lettuce with antibiotic resistant E. coli after slurry application. In J. Pulkrabová, M. Tomaniová, J. Kahl, & J. Hajslova (Eds.), *Book of Abstracts. First Int. Conference on Organic Food Quality and Health research* <http://www.fqh2011.org/>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# CONTAMINATION OF LETTUCE WITH ANTIBIOTIC RESISTANT *E. COLI* AFTER SLURRY APPLICATION

**A.N. Jensen<sup>1\*</sup>, C. Storm<sup>1</sup>, D.L. Baggesen<sup>1</sup>, A. Forslund<sup>2</sup>, and A. Dalsgaard<sup>2</sup>**

<sup>1</sup> Division of Microbiology and Risk Assessment, National Food Institute, Technical University of Denmark, Copenhagen Denmark

<sup>2</sup> Department of Veterinary Disease Biology, Faculty of Life Sciences, University of Copenhagen, Denmark

\*E-mail: anyj@food.dtu.dk; Tel: 04535886328; Fax: 04535887001

Due to disease outbreaks associated with contaminated vegetables it has been speculated to what extent this may be linked with application of animal manure as fertilizer, which is particularly practiced in organic vegetable production where conventional fertilizers are prohibited. A field survey was therefore performed to assess the survival and transfer of antibiotic-resistant *E. coli* from animal manure to lettuces, with *E. coli* serving as an indicator of bacterial enteric pathogens.

Animal slurry was applied to 3 Danish fields prior to planting of lettuce seedlings, then 5-8 weeks later at the normal time of harvest, inner and outer leaves of 10 lettuce heads were pooled into one sample unit with a total of 50 pools per field. Additionally, in one field, 15 soil samples were collected weekly until the harvest time. *E. coli* was enumerated by plating 1 mL of 10-fold serial dilutions of 5 g of homogenized sample material, i.e. manure, soil and lettuce onto Petrifilm<sup>TM</sup> Select *E. coli* count plates (3M) containing 16 mg/L streptomycin or 16 mg/L ampicillin or no antibiotics. Plates were then incubated 24 h at 44°C. Selected isolates of *E. coli* (n=83) from slurry, soil and lettuce were analysed by PFGE DNA typing for further discrimination.

The slurry applied to the fields contained 3.0-4.5 Log<sub>10</sub> *E. coli* CFU/g and resistant *E. coli* ranged from 1.0 to 4.4 Log<sub>10</sub> *E. coli* CFU/g with particular high numbers of streptomycin resistant *E. coli* in conventional pig slurry (field 1) opposed to organic cow slurry (field 2 and 3). *E. coli* was found in 36-54% of the pooled lettuce samples at the three fields with a detection limit of 10 CFU/g and 10-18% and 0-2% of pools had streptomycin and ampicillin resistant *E. coli*, respectively. Unexpectedly, the highest percentage of lettuce pools with antibiotic resistant *E. coli* were found on fields fertilized with organic cow slurry where 0.1-5% of *E. coli* was resistant opposed to 5-50% resistant *E. coli* in conventional slurry. Numbers of *E. coli* in 14-20% of pooled lettuce samples exceeded a satisfactory microbiological hygiene criteria level of 100 CFU/g. The numbers of resistant *E. coli* for both antibiotics were approximately 10-fold lower than the sensitive *E. coli*.

At the time of harvest, the numbers of *E. coli* in 5 of 15 soil samples were reduced below the detection limit and no samples exceeded 100 CFU/g, which was in contrast to the lettuce samples, where 20% of faecally contaminated samples contained >100 *E. coli*/g. This indicates that fecal contamination of crops originated from alternative sources such as contaminated water or wildlife. This was supported by genotyping of *E. coli*, where half of the 21 PFGE types were found on single occasions in either soil or lettuce, whereas the other half was found both in slurry and lettuce indicating a possible transfer.

**Key words:** lettuce, manure fertilizer, faeces contamination, resistant bacteria, food safety

**Acknowledgement:** The PathOrganic project under the CORE Organic I (Coordination of European Transnational Research in Organic Food and Farming Systems, project no. 1888) being partners of FP7 ERA-net.